



2015 AP[®] CHEMISTRY FREE-RESPONSE QUESTIONS



2. Ethene, $C_2H_4(g)$ (molar mass 28.1 g/mol), may be prepared by the dehydration of ethanol, $C_2H_5OH(g)$ (molar mass 46.1 g/mol), using a solid catalyst. A setup for the lab synthesis is shown in the diagram above. The equation for the dehydration reaction is given below.



A student added a 0.200 g sample of $C_2H_5OH(l)$ to a test tube using the setup shown above. The student heated the test tube gently with a Bunsen burner until all of the $C_2H_5OH(l)$ evaporated and gas generation stopped. When the reaction stopped, the volume of collected gas was 0.0854 L at 0.822 atm and 305 K. (The vapor pressure of water at 305 K is 35.7 torr.)



- (a) Calculate the number of moles of $C_2H_4(g)$
- that are actually produced in the experiment and measured in the gas collection tube and
 - that would be produced if the dehydration reaction went to completion.
- (b) Calculate the percent yield of $C_2H_4(g)$ in the experiment.

Because the dehydration reaction is not observed to occur at 298 K, the student claims that the reaction has an equilibrium constant less than 1.00 at 298 K.

- (c) Do the thermodynamic data for the reaction support the student's claim? Justify your answer, including a calculation of ΔG_{298}° for the reaction.

2015 AP[®] CHEMISTRY FREE-RESPONSE QUESTIONS

- (d) The Lewis electron-dot diagram for C_2H_4 is shown below in the box on the left. In the box on the right, complete the Lewis electron-dot diagram for C_2H_5OH by drawing in all of the electron pairs.



- (e) What is the approximate value of the $C-O-H$ bond angle in the ethanol molecule?
- (f) During the dehydration experiment, $C_2H_4(g)$ and unreacted $C_2H_5OH(g)$ passed through the tube into the water. The C_2H_4 was quantitatively collected as a gas, but the unreacted C_2H_5OH was not. Explain this observation in terms of the intermolecular forces between water and each of the two gases.